

(c) *Related tolerances.* See § 556.240 of this chapter.

(d) *Conditions of use.* For implantation in steers and heifers as follows:

(1) *Amount.* Insert one 25.7-mg implant every 200 days; insert one 43.9-mg implant every 400 days.

(2) *Indications for use.* For increased rate of weight gain in suckling and pastured growing steers; for improved feed efficiency and increased rate of weight gain in confined steers and heifers. No additional effectiveness may be expected from reimplanting in less than 200 days for the 25.7-mg implant or 400 days for the 43.9-mg implant.

(3) *Limitations.* For subcutaneous ear implantation in steers and heifers only. Safety and effectiveness have not been established in veal calves. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

[69 FR 67818, Nov. 22, 2004, as amended at 77 FR 31723, May 30, 2012]

§ 522.842 Estradiol benzoate and testosterone propionate.

(a) *Sponsors.* See sponsors in § 510.600(c) of this chapter for use as in paragraph (c) of this section.

(1) No. 054771 for use as in paragraph (c)(1)(i), (c)(2), and (c)(3) of this section.

(2) No. 000986 for use as in paragraph (c) of this section.

(b) *Related tolerances.* See §§ 556.240 and 556.710 of this chapter.

(c) *Conditions of use.* For implantation in heifers as follows:

(1) *Amount.* (i) 20 milligrams (mg) estradiol benzoate and 200 mg testosterone propionate (one implant consisting of 8 pellets, each pellet containing 2.5 mg estradiol benzoate and 25 mg testosterone propionate) per implant dose.

(ii) 20 mg estradiol benzoate and 200 mg testosterone propionate (one implant consisting of 9 pellets, each of 8 pellets containing 2.5 mg estradiol benzoate and 25 mg testosterone propionate, and 1 pellet containing 29 mg tylosin tartrate) per implant dose.

(2) *Indications for use.* For increased rate of weight gain and improved feed efficiency.

(3) *Limitations.* For heifers weighing 400 pounds or more; for subcutaneous

ear implantation, one dose per animal; not for use in dairy or beef replacement heifers. Safety and effectiveness have not been established in veal calves. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

[69 FR 68252, Nov. 24, 2004, as amended at 77 FR 31723, May 30, 2012; 79 FR 16187, Mar. 25, 2014]

§ 522.850 Estradiol valerate and norgestomet in combination.

(a) *Specifications.* The product is a two-component drug consisting of the following:

(1) An implant containing 6.0 milligrams of norgestomet.

(2) An injectable solution (sesame oil) containing 3.0 milligrams of norgestomet and 5.0 milligrams of estradiol valerate per 2 milliliters.

(b) *Sponsor.* See 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* One implant and 2 milliliters of injection at time of implantation.

(2) *Indications for use.* For synchronization of estrus/ovulation in cycling beef cattle and non-lactating dairy heifers.

(3) *Limitations.* Insert implant subcutaneously in the ear only; then immediately inject solution intramuscularly only. Counting the day of implantation as day 1, remove the implant on day 10. Collect all implants as they are removed and burn them. While animals are restrained for artificial insemination, avoid other treatments such as vaccinations, dipping, pour-on grub and louse prevention, spraying, etc. When inseminating without estrus detection, the entire treated group should be started at 48 hours after the last implant has been removed and should be completed within 6 hours. Where estrus detection is preferred, insemination should be approximately 12 hours after first detection of estrus. Those that do not conceive can be re-bred when they return to estrus approximately 17 to 25 days after implant removal. Do not use in